

A Hierarchical Bayesian Approach for Aerosol Retrieval Using MISR Data

Yueqing Wang ^{1,*}, Xin Jiang ^{2,†}, Bin Yu ^{1,3}, Ming Jiang ^{2,4}

July 2011

¹ Department of Statistics, University of California at Berkeley, CA 94720-3860, U.S.

² LMAM, School of Mathematical Sciences, Peking University, Beijing 100871, China.

³ Department of Electrical Engineering and Computer Sciences,
University of California at Berkeley, CA 94720-3860, U.S.

⁴ Beijing International Center for Mathematical Research, Beijing, 100871, China.

Abstract

Atmospheric aerosols can cause serious damage to human health and life expectancy. Using the radiances observed by NASA's Multi-angle Imaging SpectroRadiometer (MISR), the current MISR operational algorithm retrieves Aerosol Optical Depth (AOD) at a spatial resolution of $17.6 \times 17.6 \text{ km}^2$. A systematic study of aerosols and their impact on public health, especially in highly-populated urban areas, requires a finer-resolution estimate of the spatial distribution of AOD values.

*Corresponding author: yqwang@stat.berkeley.edu.

†Xin Jiang is now working at Netease Youdao. 1 Zhongguancun East Road, Haidian District, Beijing, 100084 China.

We embed MISR’s operational weighted least squares criterion and its forward simulations for AOD retrieval in a likelihood framework and further expand it into a Bayesian hierarchical model to adapt to a finer spatial scale of $4.4 \times 4.4 \text{ km}^2$. To take advantage of AOD’s spatial smoothness, our method borrows strength from data at neighboring pixels by postulating a Gaussian Markov Random Field prior for AOD. Our model considers both AOD and aerosol mixing vectors as continuous variables. The inference of AOD and mixing vectors is carried out using Metropolis-within-Gibbs sampling methods. Retrieval uncertainties are quantified by posterior variabilities. We also implement a parallel MCMC algorithm to reduce computational cost. We assess our retrievals performance using ground-based measurements from the AErosol RObotic NETwork (AERONET), a hand-held sunphotometer and satellite images from Google Earth.

Based on case studies in the greater Beijing area, China, we show that a 4.4 km resolution can improve the accuracy and coverage of remotely-sensed aerosol retrievals, as well as our understanding of the spatial and seasonal behaviors of aerosols. This improvement is particularly important during high-AOD events, which often indicate severe air pollution.

KEYWORDS: Hierarchical Bayesian model, MCMC, spatial dependence, fine retrieval resolution, remote sensing

1 Motivation

Atmospheric aerosols, a complex mixture of solid particles and liquid droplets in the air, can significantly affect human health and life expectancy. Studies by Monleau *et al.* and Pope *et al.* show that when inhaled, aerosols can penetrate cell membranes, then migrate and seriously damage human respiratory and cardiovascular systems [22] and the brain [21]. Short-term impacts include irritation to eyes, nose and throat; upper respiratory

infections including pneumonia and bronchitis; and stroke or death from cardiovascular causes. The “Great Smog” in London in 1952, caused by burning coal for heating, left approximately 12,000 dead [1]. Continual exposure to hazardous aerosols can aggravate or complicate medical conditions in the elderly [4]. Aerosols from silica and diesel can cause serious diseases including silicosis and black lung. In addition, aerosols with an aerodynamic diameter less than $2.5\ \mu\text{m}$, such as black carbon, can severely reduce ground-level visibility. Profiling the spatial distribution of aerosols at a fine resolution is critical for air quality and public health studies, especially in urban areas with complex anthropogenic aerosol sources, such as vehicles, power plants, and factories that burn fossil fuels.

There are two approaches to measure the spatial distribution of aerosols: through ground-based measurements and remote-sensed radiance imageries. Both adopt the same quantification of aerosols, the spectral Aerosol Optical Depth (AOD), the negative logarithm of the fraction of radiation not scattered or absorbed on a path¹. AOD at different spectral bands can be viewed as known functions of AOD at the green band using the Angström exponent [20]. For notational simplicity, we refer to AOD throughout this paper as that at the green band. With either approach, the spatial and temporal variabilities of aerosols require continual observations and computationally efficient analyses.

The AErosol RObotic NETwork (AERONET) [13] provides a data archive of local AOD values using a network of automatic sun photometers (Figure 1, left panel) located at more than 400 stations on the Earth’s surface. It measures AOD from every half hour to every two hours, with AOD uncertainties $< \pm 0.01$ at wavelengths $> 440\ \text{nm}$ [16]. AERONET measurements are widely accepted as the gold standard to validate AOD estimates from other data sources. The sparse and heterogeneous locations of AERONET stations, however, make it difficult to directly use their measurements to study AOD’s spatial behaviors.

Remote-sensing radiometers offer a better spatial coverage by retrieving AOD from radiance imageries over the Earth’s entire surface: the Multi-angle Imaging SpectroRa-

¹For example, an AOD value of 2.5 corresponds to 92% of radiation scattered or absorbed.




Figure 1: AERONET sun photometer at Avignon, France (left) and MISR cameras (right).

diometer (MISR) aboard the NASA Earth Observing System Terra satellite (Figure 1, right panel). MISR views the daylit Earth atmosphere almost simultaneously at nine angles along its track. It provides a four-spectral imagery² at a $1.1 \times 1.1 \text{ km}^2$ resolution for all spectral bands, but a $275 \times 275 \text{ m}^2$ resolution for the red.

To quantitatively represent aerosol mixtures, aerosol particles are characterized and categorized according to their properties such as radius and single scattering albedo (SSA)³. A category is referred to as a component aerosol. Then an aerosol mixture is identified by its composition: a collection of M component aerosols and their mixing vector relative to the M components. Elements of the M -dimensional mixing vector add up to 1, indicating mixing percentages of the M components. To simplify remote-sensing retrieval, MISR operational algorithm considers only 21 component aerosols and 74 pre-fixed compositions⁴.

Based on these 74 compositions and a discrete grid of AOD values, forward radiative transfer (RT) calculations simulate radiances in 36 channels (9 viewing angles $\times$ 4 spectral bands). The results form the Simulated MISR Ancillary Radiative Transfer (SMART) Dataset. The MISR operational aerosol retrieval algorithm compares SMART data to the

²The four spectral bands are blue, green, red, and near-infrared. More details of MISR observations are included in Appendix 7.1.

³SSA is defined as the ratio of scattered radiation to total extinct radiation (scattered and absorbed).

⁴Examples of MISR's 21 component aerosols and 74 compositions are displayed in Table 1 and Table 2 in Appendix 7.2. The non-zero elements of these composition vectors are no more than three.

MISR observations using a weighted least squares criterion to determine whether they provide good fits to the data, i.e., a “successful” retrieval⁵.

MISR’s unique design of multiple viewing angles provides an enhanced sensitivity to aerosol scattering and cloud reflective effects [8], giving MISR a significant advantage over other remote-sensing instruments. Validated by AERONET measurements, MISR field instruments, and airplane campaigns [5], MISR’s retrievals have shown to be informative in characterizing aerosols. Previous studies include those on wildfire smoke [18], mineral dusts [19], and climate changing aerosols [24].

MISR’s ability to capture aerosol-related information make it well suited to assist studies on aerosols’ impact on public health. MISR operational retrievals at the 17.6 km resolution, however, are too coarse to determine the relationship between aerosols and human health. The heterogeneity of urban aerosols within an area of $17.6 \times 17.6 \text{ km}^2$ makes a finer resolution desirable. For example, San Francisco is represented by less than half of a pixel on MISR output imagery. Yet the residents of San Francisco are exposed to varying level of air pollution. One also needs to add more varieties beyond the 74 pre-fixed compositions to capture the growing heterogeneity of urban aerosols.

Finer-resolution retrievals with greater varieties of aerosol compositions lead to a larger number of parameters for us to estimate. This retrieval is possible if we take advantage of AOD’s spatial smoothness and reduce the collection of 21 component aerosols to a smaller subset, say four. We chose to take a MISR Block⁶ as a data unit to balance the coverage of a greater metropolitan area and computational cost.

Our study proposes a Bayesian hierarchical model to retrieve AOD values and mixing vectors given a collection of four component aerosols at a resolution of $4.4 \times 4.4 \text{ km}^2$. The retrievals are based on MISR observations. The four component aerosols are chosen according to current knowledge of aerosol conditions for the study region. We adopt a

⁵More details on MISR operational aerosol retrieval algorithm are included in Appendix 7.2.

⁶MISR observes the Earth’s surface in 233 swaths; each swath contains 180 $560 \times 140 \text{ km}^2$ MISR Blocks.

likelihood framework based on MISR’s weighted least squares and construct a Bayesian hierarchy to incorporate AOD’s spatial smoothness into the model. In particular, we introduce a Gaussian Markov Random Field (GMRF) prior to model AOD’s spatial dependence structure. The movement and dispersion of air particles in the atmosphere justify this spatial smoothness of AOD from a physical viewpoint. To flexibly describe various aerosol conditions, our model regards AOD values and mixing vectors as continuous parameters. This expands the variety of possible compositions beyond the 74 MISR-designated choices. We show how this enriched variety is necessary for retrieving heterogeneous urban aerosols. We have not introduced an explicit spatial dependence structure for the mixing vectors of pixels in one MISR Block. Instead, the spatial smoothness of mixing vectors is implicitly enforced by the smoothness of AOD values through the hierarchy. We introduce conjugate hyperpriors to determine the parameters in the priors.

The posterior inference of AOD and mixing vectors is carried out using Markov Chain Monte Carlo (MCMC) sampling methods, particularly Metropolis-within-Gibbs. The sampling methods also allow us to quantify the retrieval uncertainties by posterior variability. The algorithm, however, is computationally intense. We develop a parallel MCMC algorithm by breaking the MISR block into smaller patches to enable parallel samplings while maintaining the overall smoothness using summary statistics. We show that retrievals from the two algorithms are consistent, with an increase in computational speed for the latter.

To assess the performance of our methods, we apply them to retrieve AOD and mixing vectors for the greater Beijing area in China. Our retrievals are tested against ground-based measurements from two AERONET stations in the area. They show improvement on accuracy and retrieval coverage, especially in high-AOD events. We also include geographic information from Google Earth to qualitatively validate our retrievals.

The rest of the paper is organized as follows: Section 2 provides the rationale and details of our Bayesian model for retrieving AOD values and mixing vectors, while Section 3

details our MCMC algorithms. Section 4 contains case studies for model validation and interpretation, comparing our results with MISR’s retrievals and AERONET measurements. In particular, Section 4.4 uses Google Earth to investigate locations with extreme AOD values, relating the retrievals to local geophysical conditions and levels of anthropogenic activities. Section 4.5 illustrates the necessity to include a richer variety of compositions. Section 5 summarizes the results and suggests directions for future research.

2 Hierarchical Bayesian Model

Our study objective is to establish a more detailed data-driven description of the relationships among radiances, AOD, and aerosol compositions to assist aerosol-related health studies. The MISR operational retrieval algorithm provides this information by comparing the observed and RT calculated radiances, but it is limited within the 74 pre-fixed compositions and a discrete grid of AOD values. We propose to understand a greater variety of optical behaviors of aerosols by considering AOD values and mixing vectors as continuous variables, given a fixed collection of four component aerosols. For the greater Beijing area, the collection includes spherical non-absorbing aerosols without sulfate, spherical non-absorbing aerosols with sulfate, spherical absorbing aerosols, and grains (dust).

Each MISR Block contains 256 pixels ($8 \text{ rows} \times 32 \text{ columns}$) at the 17.6 km resolution in MISR retrievals. The number of pixels rises to 4,096 ($32 \text{ rows} \times 128 \text{ columns}$) as we retrieve at 4.4 km resolution, presenting a more complex problem with approximately 16,384 parameters to estimate for each of 41,940 MISR Blocks. On the other hand, air particles interact in the atmosphere within a certain range; consequently, they affect aerosol conditions in near neighborhoods [2] [26]. This suggests a stronger spatial dependence among adjacent pixels at a finer scale. When modeling at a fine resolution, therefore, it is beneficial and necessary to borrow strength from AOD’s spatial smoothness to reduce model complexity. In particular, we construct a hierarchical Bayesian model with a built-in

spatial dependence structure using a GMRF prior for AOD values.

In Section 2.1, we introduce the notations and define the likelihood function, based on MISR's weighted least squares criterion. Section 2.2 introduces our Bayesian hierarchy. Unless specified otherwise, our Bayesian retrievals are at a 4.4 km resolution.

2.1 Defining the Likelihood Function

Let $p = 1, \dots, P$ denote the $P = 4,096$ pixels on a two-dimensional lattice in a MISR Block at 4.4 km. For pixel p , MISR observes the top-of-atmosphere radiances $\mathbf{L}_p = (L_{1p}, \dots, L_{Cp})$ at MISR's $C = 36$ channels. Every channel c is weighted by $\frac{1}{2\sigma_c^2}$, $c = 1, \dots, 36$, to account for different measurement errors. For pixel p , let $\tau_p \in \mathbf{R}$ denote its AOD value, and $\boldsymbol{\theta}_p = (\theta_{p1}, \dots, \theta_{pM}) \in \mathbf{R}^M$ the mixing vector of the M component aerosols involved. $\sum_{m=1}^M \theta_{pm} = 1$. Let $\mathbf{L} = \{\mathbf{L}_1, \dots, \mathbf{L}_P\}$, $\boldsymbol{\tau} = \{\tau_1, \dots, \tau_P\}$, and $\boldsymbol{\theta} = \{\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_P\}$. For each pixel p independently, the RT calculated radiances $\mathbf{L}^{RT} = (L_1^{RT}, \dots, L_C^{RT})$ can be viewed as functions of $(\tau_p, \boldsymbol{\theta}_p)$, relative to the collection of component aerosols involved. The MISR operational retrieval algorithm exhaustively searches over combinations of pre-fixed AOD values and 74 compositions to match $\mathbf{L}^{RT}$ to the observed, based on the following weighted least squares [6]:

$$\chi_p^2 = \sum_{c=1}^C \frac{(L_{cp} - L_c^{RT}(\tau_p, \boldsymbol{\theta}_p))^2}{2\sigma_c^2}. \quad (1)$$

The combinations of AOD and compositions whose χ_p^2 satisfy a pre-established threshold are considered successful and the averages of all successful retrievals are the outputs.

In this paper, we turn the weighted least squares in (1) into a likelihood function to assess how probable are the observations $\mathbf{L}$ given $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$. We quantify this probability through a Gaussian density based on the Euclidean distance between $\mathbf{L}$ and $\mathbf{L}^{RT}$. Aerosol retrievals constrained by (1) can be formalized as inference of $(\boldsymbol{\tau}, \boldsymbol{\theta})$ based on the following

likelihood function,

$$p(\mathbf{L}|\boldsymbol{\tau}, \boldsymbol{\theta}) \propto \exp \left\{ - \sum_{c=1}^C \sum_{p=1}^P \frac{(L_{cp} - L_c^{RT}(\boldsymbol{\tau}_p, \boldsymbol{\theta}_p))^2}{2\sigma_c^2} \right\}. \quad (2)$$

In the MISR operational retrieval, $\{\sigma_c^2\}_{c=1}^C$ are explicitly assigned as the minimum of 0.04 and $\bar{L}_c = (\sum_{p=1}^P L_{cp})/P$ (Appendix 7.2). In our model, $\{\sigma_c^2\}_{c=1}^C$ are considered unknown parameters and are to be estimated together with $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$. We set the number of component aerosols $M = 4$ and case studies show the sufficiency of this choice in our cases.

We adopt a Bayesian approach since it easily incorporates smoothness in $\boldsymbol{\tau}$ through their prior distributions. It also provides uncertainty measures through posterior variances. In the following section, we describe our hierarchical model by assigning reasonable priors to the unobserved variables and building conditional relationships within the Bayesian hierarchy.

2.2 Construction of the Priors and Conditional Probabilities

For fixed atmospheric pressures, humidity and wind levels, the top-of-atmosphere radiances $\mathbf{L}$ are mainly determined by $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$. We use the hierarchy's first level to indicate the dependence of $\mathbf{L}$ on $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$. Prior distributions for $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$ are further assumed to be independent, i.e., $p(\boldsymbol{\tau}, \boldsymbol{\theta}) = p(\boldsymbol{\tau})p(\boldsymbol{\theta})$ to simplify computation. The posterior distribution of $\boldsymbol{\tau}$ will influence that of $\boldsymbol{\theta}$ by interacting within the hierarchy, and vice versa.

Priors for $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$ are postulated to capture their smoothness calibrated by hyperparameters κ and α respectively. A GMRF prior is adopted to describe $\boldsymbol{\tau}$'s spatial dependence. Though there is no explicit spatial structure for $\boldsymbol{\theta}$, the interactions between $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$ through the hierarchy implicitly enforce a spatial dependence for $\boldsymbol{\theta}$. Figure 2 provides a graphical representation of the hierarchy. It characterizes the dependence of $\mathbf{L}$ on parameters $\boldsymbol{\tau}$ and $\boldsymbol{\theta}$ and the independence among the hyperparameters. The inference of parameters and hyperparameters, using MCMC sampling methods, is discussed in Section 3.




Figure 2: Graphical representation of the hierarchical Bayesian model. Green square: observations; red circle: model parameters; blue circle: model hyperparameters.

2.2.1 Prior Beliefs about AOD's Spatial Dependence

We model the spatial dependence structure of AOD values using an intrinsic GMRF prior of first order [23]. Define κ as the homogenous scaler precision and use $\sim$ to indicate spatial adjacency. The following prior is invariant to perturbation by the same constant to τ of all pixels [3],

$$p(\tau|\kappa) \propto \kappa^{\frac{p-1}{2}} \exp \left\{ -\frac{\kappa}{2} \sum_{p' \sim p} (\tau_{p'} - \tau_p)^2 \right\}. \quad (3)$$

This allows us to model the smoothness of τ , regardless of its unknown overall level, by penalizing sharp changes in τ among adjacent pixels.

The above GMRF prior is governed by one precision parameter κ . The larger κ is, the smoother the region's AOD values are, with $\frac{1}{\sqrt{\kappa}}$ as AOD's precision. For some regions, however, a more complicated GMRF prior might be necessary. For example, a continuing pattern of wind directions might make it useful to distinguish a upwind pixel from a downwind pixel. This paper works with a homogenous precision κ for τ 's GMRF prior.

To estimate κ , we assign κ a hyperprior. Since $p(\tau|\kappa)$ belongs to an exponential family, we choose its natural conjugate as $p(\kappa)$, a noninformative Gamma distribution. The posterior is also a Gamma distribution,

$$p(\kappa|\tau) \propto \kappa^{(p-1)/2-1} \exp \left\{ -\frac{1}{2} \kappa \sum_{p': p' \sim p} (\tau_{p'} - \tau_p)^2 \right\}. \quad (4)$$

2.2.2 Prior Specification for Aerosol Compositions

Prior information on aerosol compositions is incorporated in the model through the choice of the $M = 4$ component aerosols involved, based on geophysical knowledge of the study region. To model the mixing vectors θ for these M component aerosols, we adopt an M -dimensional Dirichlet prior with parameter $\alpha = (\alpha_1, \dots, \alpha_M)$. By adjusting the magnitude of α , we can control the sparsity of the mixings of component aerosols. Conditioning on α , the mixing vectors θ_p for each pixel p are considered to be independent,

$$p(\theta|\alpha) = \prod_{p=1}^P p(\theta_p|\alpha) = \prod_{p=1}^P \frac{\Gamma(\sum_{m=1}^M \alpha_m)}{\prod_{m=1}^M \Gamma(\alpha_m)} \theta_{p1}^{\alpha_1-1} \dots \theta_{pM}^{\alpha_M-1}. \quad (5)$$

In general cases with no information on the mixing's sparsity, we assign α a hyperprior to estimate it. Since (5) belongs to an exponential family, we take its conjugate as $p(\alpha)$,

$$p(\alpha) \propto \exp \left(\sum_{m=1}^M (\alpha_m - 1) \gamma_{0,m} - \eta \left(\sum_{m=1}^M \log \Gamma(\alpha_m) - \log \Gamma \left(\sum_{m=1}^M \alpha_m \right) \right) \right).$$

η is chosen as 0 and $\{\gamma_{0,m}\}_{m=1}^M$ as 1 to obtain an exponential hyperprior. The posterior is,

$$p(\alpha|\theta) \propto \exp \left\{ \sum_{m=1}^M (\alpha_m - 1) \left(\sum_{p=1}^P \log \theta_{pm} + 1 \right) - P \times \left(\sum_{m=1}^M \log \Gamma(\alpha_m) - \log \Gamma \left(\sum_{m=1}^M \alpha_m \right) \right) \right\}.$$

2.2.3 Hyperpriors for σ

The likelihood function for $\sigma = (\sigma_1, \dots, \sigma_C)$, $p(L|\sigma, \tau, \theta)$, follows a normal distribution with known mean and unknown variance. We adopt a conjugate noninformative scaled inverse- χ^2 prior for σ to model the channel weights $\{\frac{1}{2\sigma_c^2}\}$. Thus, $p(\sigma_c^2) \propto \sigma_c^{-2}$ ($c = 1, \dots, C$), and the conditional posterior also follows the scaled inverse- χ^2 distribution,

$$p(\sigma_c^2|\tau, \theta, L) \propto (\sigma_c^2)^{-(\frac{P}{2}+1)} \exp \left\{ -\frac{\sum_{p=1}^P (L_{cp} - L_c^{RT}(\tau_p, \theta_p))^2}{2\sigma_c^2} \right\}.$$

3 MCMC Retrieval Algorithms

Based on the hierarchical Bayesian model previously developed, this section derives marginal posterior distributions of AOD values τ and mixing vectors θ and devises two MCMC algorithms to sample from the posteriors. Using MISR observed radiances as input, the sampled posterior means are used as our retrieval outputs.

3.1 Posterior Distributions of AOD and compositions

The full Bayesian model discussed above can be summarized as follows:

$$L_p | \tau_p, \theta_p \sim \mathcal{N}(L^{RT}(\tau_p, \theta_p), \sigma^2), \quad p = 1, \dots, P,$$

$$\tau | \kappa \sim \text{GMRF}(\kappa),$$

$$\theta | \alpha \sim \text{Dirichlet}(\alpha),$$

$$\sigma \sim \text{scaled inverse } -\chi^2(\nu_0),$$

$$\kappa \sim \text{Gamma}(\alpha_0, \beta_0),$$

$$p(\alpha) \sim \text{Exp}\left(\sum_{m=1}^M (1 - \alpha_m)\right).$$

With no additional information on the hyperparameters, ν_0 , α_0 , and β_0 are chosen to be 0 for convenience and later shown to be robust. The marginal posterior of τ is,

$$p(\tau | \theta, \kappa, \sigma, L) \propto \exp \left\{ -\frac{1}{2} \kappa \sum_{p': p' \sim p} (\tau_{p'} - \tau_p)^2 - \sum_{c=1}^C \sum_{p=1}^P \frac{(L_{cp} - L_c^{RT}(\tau_p, \theta_p))^2}{2\sigma_c^2} \right\}. \quad (6)$$

The marginal posterior distribution of θ can be expressed as,

$$p(\theta | \tau, \alpha, \sigma, L) \propto \exp \left\{ \sum_{p=1}^P \sum_{m=1}^M (\alpha_m - 1) \log \theta_{pm} - \sum_{c=1}^C \sum_{p=1}^P \frac{(L_{cp} - L_c^{RT}(\tau_p, \theta_p))^2}{2\sigma_c^2} \right\}. \quad (7)$$

Both posteriors contain an RT-based $\mathbf{L}^{RT}$, can be obtained at necessary values through interpolations from MISR SMART Dataset [7], using τ_p and θ_p as input (Appendix 7.4). These non-closed-form posteriors, however, are difficult to directly sample from. A Metropolis-within-Gibbs sampler is used when RT-based $\mathbf{L}^{RT}$ are involved.

3.2 Metropolis-within-Gibbs Sampling from the Posterior Distributions

The Gibbs sampler [11] is a numerical technique to sample from a joint posterior distribution, $p(\tau, \theta, \sigma, \kappa, \alpha | \mathbf{L})$ in our case. We sample for τ_p and θ_p using a Metropolis-Hastings (M-H) sampler, for each pixel p on the MISR Block column-by-column and pixel-by-pixel. The following proposal distribution is used in M-H for τ_p ,

$$p(\tau_p | \tau_{-p}) \propto \exp \left(-\frac{n_p \kappa}{2} (\tau_p - \frac{1}{n_p} \sum_{p': p' \sim p} \tau_{p'})^2 \right),$$

where n_p is the number of adjacent pixels to pixel p . A Dirichlet proposal distribution with parameter α is used for θ_p . Denote vector $(\tau_p, \dots, \tau_{p'})$ by $\tau_{p:p'}$ and similarly for θ and α . Given initializations $(\tau^{(0)}, \theta^{(0)}, \sigma^{(0)}, \kappa^{(0)}, \alpha^{(0)})$, the Gibbs sampler proceeds as follows:

Metropolis-within-Gibbs Algorithm (M-w-G)

At step t , iterate the following process:

- Use M-H to sample $\tau_1^{(t)} \sim p(\tau_1 | \tau_{2:P}^{(t-1)}, \theta^{(t-1)}, \sigma^{(t-1)}, \kappa^{(t-1)}, \mathbf{L})$;
 Use M-H to sample $\tau_2^{(t)} \sim p(\tau_2 | \tau_1^{(t)}, \tau_{3:P}^{(t-1)}, \theta^{(t-1)}, \sigma^{(t-1)}, \kappa^{(t-1)}, \mathbf{L})$; ...;
 Use M-H to sample $\tau_P^{(t)} \sim p(\tau_P | \tau_{1:(P-1)}^{(t)}, \theta^{(t-1)}, \sigma^{(t-1)}, \kappa^{(t-1)}, \mathbf{L})$.
- Use M-H to sample $\theta_1^{(t)} \sim p(\theta_1 | \tau^{(t)}, \theta_{2:P}^{(t-1)}, \sigma^{(t-1)}, \alpha^{(t-1)}, \mathbf{L})$; ...;
 Use M-H to sample $\theta_P^{(t)} \sim p(\theta_P | \tau^{(t)}, \theta_{1:(P-1)}^{(t)}, \sigma^{(t-1)}, \alpha^{(t-1)}, \mathbf{L})$.
- Use M-H to sample $\sigma_1^{(t)} \sim p(\sigma_1^2 | \tau^{(t)}, \theta^{(t)}, \sigma_{2:C}^{(t-1)}, \mathbf{L})$; ...;
 Use M-H to sample $\sigma_C^{(t)} \sim p(\sigma_C^2 | \tau^{(t)}, \theta^{(t)}, \sigma_{1:(C-1)}^{(t)}, \mathbf{L})$.
- Sample $\kappa^{(t)} \sim p(\kappa | \tau^{(t)})$.
- Use M-H to sample $\alpha_1^{(t)} \sim p(\alpha_1 | \theta^{(t)}, \alpha_{2:M}^{(t-1)})$; ...;
 Use M-H to sample $\alpha_M^{(t)} \sim p(\alpha_M | \theta^{(t)}, \alpha_{1:(M-1)}^{(t)})$.

The above cycle generates a realization of a Markov chain, which gives approximate samples from the marginal posteriors after a successful burn-in process [11]. Appendix 7.5 contains details on the stability of sampled posterior mean over different initializations.

Whenever we used the Metropolis-Hastings algorithm, we checked that the acceptance rate was roughly between 25% and 50% to secure adequate mixing of posterior samples [9]. We also used a variety of diagnostic tools to ensure the convergence of the Markov chains, including the potential scale reduction $\hat{R}$ [10] on several parallel Markov chains with different initializations⁷. We ran the chains until $\hat{R}$ is less than 1.1 or 1.2, using $\hat{R}$ of the logarithm of the posterior distribution as a benchmark. In addition, a geometric decay of the autocorrelation as a function of the lag also validated the mixing of our chains.

In the end, we estimate the posterior mean based on the samples and consider it the point estimates for τ and θ . While the MCMC algorithm enables us to handle a hierarchy whose complexity precludes fitting by analytical methods, its computation intensity limits its online use. Next, we propose a parallel MCMC algorithm to reduce computational cost.

3.3 A Parallel MCMC Algorithm

Many MCMC sampling algorithms for spatial data suffer from high computational cost caused by the large dimensionality of spatial data. On a 4.4 km scale, our MCMC algorithm simulates samples for more than 16,000 variables⁸ for one MISR Block. The large computation cost is exacerbated by the non-closed form of the posterior distributions. It is crucial, therefore, to develop an efficient and reasonably accurate algorithm.

We devise a parallel MCMC algorithm to improve the computational efficiency. In this algorithm, each MISR block is divided into 2×8 patches of equal size with at least four overlapping columns and rows between adjacent patches. The M-w-G sampler is applied to each patch independently to generate samples of τ and θ for these patches. This

⁷For example, cases are inspected with initial values of κ as 10, 100, 200, 800.

⁸Excluding cloudy pixels can sometimes reduce the total dimensions to around 5,000 for one MISR Block.

independent sampling on different patches can therefore benefit from parallel computing.

Information on AOD's spatial dependence structure is to be maintained across the entire MISR Block to achieve an accurate estimate of its spatial smoothness level. On that account, we let the parallel MCMC algorithm periodically exchange spatial smoothness information among all patches on the entire MISR Block. Given κ 's conditional posterior,

$$p(\kappa|\tau) \propto \kappa^{(P-1)/2-1} \exp\{-\frac{1}{2}\kappa \sum_{p \sim p'} (\tau_p - \tau_{p'})^2\},$$

and summary statistic, $T_\kappa = \sum_{p \sim p'} (\tau_p - \tau_{p'})^2$, it follows that $p(\kappa|\tau) = p(\kappa|T_\kappa)$. Hence T_κ is a sufficient statistic that holds all information on calibration κ for AOD's spatial smoothness across the entire MISR block. We now describe the parallel MCMC algorithm in detail:

Parallel MCMC Algorithm

1. Obtain a standard MISR block of 32×128 pixels. Divide the block into 2×8 patches, each of 20×20 pixels, with at least 4 overlapping columns/rows between adjacent patches.
2. At step t , iterate the following process:
 - Use M-w-G algorithm to sample $\tau \sim p(\tau|\theta, \kappa, \sigma, L)$, $\theta \sim p(\theta|\tau, \alpha, \sigma, L)$, $\sigma \sim p(\sigma|T_\sigma^{(t)})$, $\kappa \sim p(\kappa|T_\kappa^{(t)})$, $\alpha \sim p(\alpha|T_\alpha^{(t)})$ within each patch in parallel for 50 iterations.
 - Average the samples of the overlapping pixels between any two adjacent patches.
 - Calculate summary statistics using current samples,

$$T_{\sigma_c}^{(t+1)} = \sum_{p=1}^P (L_{cp} - L_c^{RT}(\tau_p, \theta_p))^2, c = 1, \dots, C,$$

$$T_\kappa^{(t+1)} = \sum_{p \sim p'} (\tau_p - \tau_{p'})^2,$$

$$T_{\alpha_m}^{(t+1)} = \sum_{p=1}^P \log \theta_{pm}, m = 1, \dots, M.$$

Given that the hyperparameters control the level of spatial smoothness of model parameters in all patches, the spatially dependence structure of this entire imagery is preserved, while the parallel-patch samplings improve the algorithm's computation efficiency. This parallel MCMC sampling scheme can be generalized for other computations on spatial data using sufficient statistics to maintain a global spatial structure. The above process can

be automated using the Perl programming language. For one MISR block at a resolution of 4.4 km, the computational time for the parallel MCMC algorithm is less than one-fifth of that for the global MCMC sampling algorithm (Appendix 7.5).

The global and the parallel MCMC algorithms produce reasonably consistent results. Figure 3 shows two scatterplots of AOD retrievals from global and parallel MCMC algorithms after the same number of iterations. The outputs generally agree, except for a small group of pixels that mostly lie on the edges of patches. The spatial smoothness is interrupted between patches; therefore, the benefits of a stabilizing factor from neighboring pixels are lost. This confirms the importance of maintaining an appropriate spatial dependence structure. Increasing the number of iterations and communications of summary statistics, as well as moving the edges of patches, reduces the disagreement.

The next section evaluates the performance of our Bayesian retrievals' using case studies. For non-operational model validations, we apply the global MCMC algorithm to avoid inconsistency in number of iterations for different retrievals.




Figure 3: Two examples of the two MCMC algorithms. x: AOD retrievals from the global MCMC sampling algorithm; y: AOD retrievals from the parallel MCMC algorithm.

4 Assessing MCMC Algorithm Performance: Case Studies on Aerosol Retrievals for the Greater Beijing Area, China

In this section, comparison studies to MISR outputs are presented for the greater Beijing area and their differences discussed. We validate our AOD retrievals using AERONET measurements and satellite images from Google Earth. Through case studies, we demonstrate the importance of high-resolution retrieval to gain more information on aerosols, and a greater variety of compositions to improve retrieval accuracy and coverage. All retrievals in this section are based on MISR observations over the greater Beijing area (latitude 38.95N~40.15N; longitude: 115.57E~119.50E).

4.1 Historical and Present Background of Air Pollution in Beijing

There are two main reasons for high AOD values in Beijing. First, anthropogenic activities produce various air pollutants such as waste gas from over two million automobiles, industrial emissions, and dust from over 5,000 construction sites. Coal-dominated energy supplies burn off over 28 million tons of coal per year. Due to the presence of multiple air pollutants, Beijing bears high AOD and heterogeneous aerosol compositions. Second, geographical characteristics of the region also contribute to the buildup of aerosols. Mountains and hills surround the greater Beijing area on the east, west, and north. Aerosols accumulate until removed by rain or dispersed by wind into the plains on the south. Wind in any other direction is blocked and venting of aerosols highly restricted.

The anthropogenic behaviors and seasonal weather patterns lead to a yearly trend of air quality in Beijing. In winter, starting from mid-November, the use of central heating systems bring large amounts of sulfur dioxide, nitrogen dioxide, and other types of incompletely burned waste into the air. Enormous boilers of the heating systems generate masses of water vapor that help to increase the absorption of particles in the air. In December, the cold airflow from Siberia generates northwestern winds in Beijing that clear the accumu-

lated pollutants, leading to the lowest aerosol loadings in January. The loadings vary due to the discontinuous cold airflow. In spring, wind from Siberia and Loess Plateau of North-western China brings sandstorms. Traffic and industrial activities revive as temperatures warm up. From June to August, aerosol loadings reach the highest of the year [25]. From September to December, precipitation disperses and dissolves air pollutants [14].


Significant measures have been introduced to control air pollution in Beijing [15], but rapid economic and population growth, as well as the continuous expansion of the transportation system, have hindered their success. On March 20, 2010, Beijing experienced its biggest sandstorm since 2006. The storm originated from Inner Mongolia and haunted Beijing for over 5 hours. Characterizing AOD's spatial distribution under such extreme air conditions is especially important to understand its influence on public health.

4.2 Comparison with MISR Retrievals

To retrieve AOD, we average all possible AOD values weighted by the posterior distribution, while MISR averages successful retrievals with equal weights. This suggests consistency between our retrievals and those of MISR, and possible differences.

For a first impression of the aerosol conditions in this region on Jan 10, 2009, Figure 4 panel (a) displays the observed MISR imagery by one of the forward cameras (CF). MISR AOD retrievals are plotted at their 17.6 km resolution in panel (b). Our Bayesian AOD retrievals plotted in panel (c) at 4.4 km. Figure 4 suggests shared information in MISR and our MCMC retrievals, such as the coastline on the right, the overall AOD level, and its spatial patterns. This consistency is confirmed by the scatterplots of MISR outputs and Bayesian AOD retrievals aggregated to a 17.6 km resolution (Figure 5, left panel).


The increased diversity in our Bayesian AOD retrievals across the MISR Block (Figure 5, right panel) indicates that more details are detected as the spatial resolution improves. This is expected, since a finer resolution leads to more information observed and piped



(a) MISR radiance imagery by the nadir camera on Jan 10, 2009.



(b) MISR AOD retrievals at 17.6 km resolution.



(c) Bayesian AOD retrievals using MCMC at 4.4 km resolution.

Figure 4: AOD estimates from MISR and our Bayesian retrievals.

into the model. But the reliability of such diversity needs to be further validated by other independent sources such as ground-based measurements (Section 3.4).

The black pixels represent missing retrievals due to MISR instrumental design or masking criteria such as cloud screening. Aerosol retrievals are not attempted when clouds are detected. MISR averages the $1.1 \times 1.1 \text{ km}^2$ observations into a pixel of $17.6 \times 17.6 \text{ km}^2$ and ignores small clouds, given the cloudless areas are more than $\frac{1}{16}$ of the pixel. Small clouds negligible on the 17.6 km scale might be significant on a 4.4 km scale, so we tend to have more missing retrievals in some areas intrinsically determined by the observations. However, when none of the 74 MISR-designated compositions produce successful retrievals, MISR operational algorithm marks it as missing. Our Bayesian retrievals, with a richer variety of compositions, eliminate these unnecessary missing retrievals (Section 4.5).

To objectively evaluate the performance of our MCMC retrieval algorithms, the next section conducts a validation study using ground-based measurements.




Figure 5: Scatterplots of MISR against MCMC retrievals at an aggregated 17.6 km resolution (left, r.m.s. = 0.0295) and a 4.4 km resolution (right, r.m.s. = 0.0309).

4.3 Model Validation for Bayesian Retrievals by Ground-based Measurements

Ground-based measurements are collected at AERONET Beijing and AERONET Xianghe stations, as well as via a hand-held MICROTOPS II Sunphotometer at several locations in urban Beijing. The fixed locations of the AERONET stations and the limited travel range of the Sunphotometer’s human operator make it impossible to validate retrievals of all pixels on the MISR Block. Instead, we focus on the pixels that contain the AERONET stations and our Sunphotometer-visited locations. To match the Bayesian retrievals at 550 nm, we first convert AERONET data to a wavelength of 550 nm using AERONET estimates of Angström exponent and a linear transformation. We then average the measurements within a one-hour window when Terra carrying MISR passes over the AERONET stations. Jiang, *et al.* [17] show that a narrower time window better captures the correlations between AERONET measurements and MISR retrievals. The frequent cloudy weather in the greater Beijing area and its latitude⁹ contribute to the difficulty of matching remote-sensed versus ground-based data pairs for model validation.

⁹The Beijing city is visited by the Terra satellite every five to nine days.

4.3.1 Retrieval Validation at AERONET Beijing Station

Figure 6 shows a boxplot of our Bayesian AOD retrievals for the pixel which contains the AERONET Beijing Station¹⁰, with estimated uncertainties indicated by the box edges for inter-quartile ranges of posteriors and the whiskers drawn to the 5th and 95th percentiles. The three retrievals on March 15, April 30, and May 16, 2009 are plotted separately in the right panel to keep an appropriate scale for the left panel.

As long as a pixel is cloudless, our MCMC algorithms provide an AOD retrieval. However, the MISR operational retrieval algorithm produces missing values for 24% of the 21 cases in Figure 6. This results from the limited choices of aerosol compositions in MISR retrievals and the increasingly heterogeneous aerosol conditions in Beijing.

In general, our Bayesian retrievals show improvement in accuracy. Detailed information on aerosols are revealed by the fine-resolution retrievals. In the coarse-resolution retrievals, however, high AOD values are averaged down by its neighbors and low AOD values averaged up, resulting in the loss of useful information. The three high AOD values in the right panel of Figure 6 indicate Beijing's extreme air conditions, corresponding to 86%, 71%, and 81% reduced radiation by aerosols. For example, records of news on the




Figure 6: Validation of our AOD retrievals by measurements at AERONET Beijing Station.

¹⁰Latitude: 39.97689° North; longitude: 116.38137° East.



Figure 7: Left: air conditions in Beijing, China, on March 15, 2009; right: locations of Hebei Province and Shenyang.

Internet indicate that on March 15, 2009, the city was trapped in a sandstorm originated in Inner Mongolia (Figure 7).

4.3.2 Retrieval Validation at AERONET Xianghe Station

Figure 8 compares the remote-sensed retrievals to AERONET measurements at the pixel that contains the AERONET Xianghe station¹¹. From December to February, AERONET measurements are mostly higher than remote-sensed retrievals, but no pattern afterwards.

AERONET Xianghe station has the Jingshen Expressway to its north, which is a major path connecting two hub cities: Beijing and Shenyang¹². The northwest wind in winter carries car exhaust to the AERONET Xianghe station, possibly leading to high AOD measurements. Yet for remote-sensing retrievals, the green fields in a larger neighborhood balance this factor, which possibly results in a washed-out signal. However, the fine-resolution retrievals seem to suffer less from the balancing factors and display a better accuracy.

We would like to discuss one of the cases where our AOD retrieval is much higher than AERONET measurement: the first data point in Figure 8 (December 25, 2008). MISR operational retrieval algorithm produced no output for this day. To the east of Xianghe

¹¹Latitude: 39.75360° North; longitude: 116.96150° East.

¹²The capital and largest city of Liaoning Province in Northeast China.




Figure 8: Validation of our AOD retrieval results by AERONET measurements in Xianghe.

station in Hebei Province lie several major malls for furniture exhibition and manufacture. On December 25, 2008, the furniture companies started renovating their exhibition halls. The construction could have caused localized aerosol loadings not observed by the Xianghe AERONET site 2 km away upwind within one day, but detected by the MISR instrument and captured by our retrievals.

4.4 Validation Using Google Earth

Model validation by ground measurements is reassuring, except for the limited number of matching data pairs. For areas where ground measurements are not available, we investigate using satellite images from Google Earth.

Figure 9 shows a projection of our retrievals onto Google Earth. We take advantage of the interactive feature of Google Earth to zoom in on the map to investigate the geographical and anthropogenic conditions of each region, for example, the existence of heavy industries and transportation patterns. In particular, we examine pixels where the regional AOD values are not very smooth. Information carried by these pixels are likely to be revealed only at a fine resolution. We are able to identify possible emission sources around several locations with high AOD values as marked in Figure 9:




Figure 9: Retrieval results projected on Google Earth.

- A) A hub of several major highways, including Jingshen and Jingtang Highways;
- B) Industrialized areas, with shadows of high chimneys;
- C) Construction site pollution: building a new holiday resort.

Similarly, our investigation found locations with a low level of AOD values reveals that they have more natural features and less traffic and industries, for example:

- D) The Olympic Park;
- E) Mountain Xi, a famous hiking destination; and
- F) Beidaihe, a famous beach resort.

Google Earth, with its interactive character, can be employed to understand the behaviors of aerosols and identify locations with high or low AOD values.

4.5 Case Study for Including a Richer Variety of Aerosol Compositions

This section emphasizes the necessity of expanding MISR-designated 74 aerosol compositions. This expansion improves coverage of the MISR operational retrievals and detects

more features of aerosol behaviors (e.g., seasonal trend for certain component aerosols).




Figure 10: Case study of AOD retrievals on March 15, 2009.

For an example of the improvement on retrieval coverage, we examine March 15, 2009. MISR failed to retrieve AOD for the majority of the Block (Figure 10). Our Bayesian retrievals provide better coverage and the retrievals give distinctly high AOD values with a clear path of aerosols migrating from west to east and into the ocean. This unusual discrepancy leads us to run through the weather records: on that day, the area suffered from a sandstorm originated in Inner Mongolia, which later passed into eastern China. For areas like Beijing, which experience occasional sandstorms, the limited compositions containing grains(dust) among MISR's 74 choices could easily result in a low coverage of MISR retrievals. Similar situations might exist for other locations with usual aerosol conditions.

The retrieved mixing vectors also contains information on the regional aerosol composition and can be used to identify pollution type and source. Figure 11 shows the retrieved mixing vectors on January 10, 2009, on the left, and those on April 14, 2009, on the right. Component No.6 with sulfate tends to dominate the composition in winter due to coal




Figure 11: Retrievals of aerosol composition with $\theta = (\theta_1, \theta_2, \theta_3, \theta_4)$ plotted as lengths of segments in a column. Left: January 10, 2009; Right: April 14, 2009.

burning for heating, with No.19, grains (dust) dominating in spring due to sandstorms.

Figure 12 shows the mixing percentages of component No.19 over December 2008 to June 2009, at four different locations: the AERONET Beijing station, the AERONET Xianghe station, location (A) and (F) marked in Figure 9. For AERONET Beijing station, the percentage of grains(dust) only rose in the spring, due to the sandstorms, while the constructions around AERONET Xianghe might have raised the percentage earlier in the year. Location (A), where major highways intersect, showed a high amount of dust in its aerosol compositions through the warm seasons when traffic typically increases. The mixing percentage of No.19 at Beidaihe Resort moved relatively in consistence with AERONET Beijing station. We hope to explore this trend in future research.

In general, by correctly identifying the major pollutants for each season, we can better understand the transitions of aerosols and, therefore, take efforts to improve air quality in a more specific and to the point manner. For accuracy and coverage, it is necessary to expand the MISR-designated 74 compositions to a richer variety.




Figure 12: Mixing percentages of component No.19 from winter to spring.

5 Discussion

Aerosols serve as an important factor in climate systems and air quality. Global climate and regional wind patterns, as well as anthropogenic activities, shape aerosols into an increasingly heterogeneous system. Such complexity calls for flexible models with a compatibly fine resolution. In this paper, we have presented a Bayesian hierarchical model to retrieve AOD and mixing vectors relative to a collection of four component aerosols at a spatial resolution of 4.4 km using MISR observations. The model incorporates a spatial dependence structure to gain strength from AOD's smoothness, at the same time allows for a richer variety of aerosol mixing vectors. A more detailed AOD spatial profile is provided then validated by AERONET and Google Earth; a better retrieval coverage is obtained due to the flexible choices of aerosol compositions and improved accuracy. This improvement is particularly important during high-AOD events, which often indicate severe air pollution.

We hope that a more detailed aerosol profile will eventually expand the potential of MISR observations to assist urban air quality monitoring and public health studies. From the cases, we become more aware of the complexity in modeling aerosol conditions. External data such as Google Earth can introduce new access to model validation.

For future research, we plan to incorporate more prior information into the model, such

as wind measurements. If this information is available on an appropriate scale, it may improve retrieval performance. Our ultimate goal is to explore the possibility of using our AOD retrievals at a fine resolution to help understand the impact of AOD on public health.

6 Acknowledgments

This research is supported in part by the National Science Foundation Grants SES-0835531 (CDI), DMS-0907632, CCF-0939370, ARO Grant W911NF-11-1-0114, the National Science Foundation of China (60325101, 60872078), Key Laboratory of Machine Perception (Ministry of Education) of Peking University, and Microsoft Research of Asia. The authors would like to thank Dr. Susan Paradise, Dr. Amy Braverman, and the MISR team for their great support, Derek Bean for editing advice. We also thank the AERONET PIs, Hong-Bin Chen, Philippe Goloub, Pucai Wang, Zhanqing Li, Brent Holben, and Xiangao Xia for establishing and maintaining the Beijing and Xianghe AERONET stations, especially Pei Wang from Peking University for collecting AOD data in Beijing. We thank Graham Shapiro for implementing the projection of AOD retrievals to Google Earth. We thank Dr. Chengcai Li from Peking University and Yang Liu from Emory University for discussions.

7 Appendix

7.1 The MISR Instrument and Observations

MISR views the daylight Earth atmosphere almost simultaneously at nine angles along its track. It provides a four-spectral imagery at a $275 \times 275 \text{ m}^2$ spatial resolution. The radiances in all spectral bands except for the red are aggregated onboard to a $1.1 \times 1.1 \text{ km}^2$ resolution to reduce data transmission from Terra. The MISR cameras cover the Earth's surface in 233 geographically distinct but overlapping paths of approximately 560 km wide, extending from the Arctic to Antarctica. Every swath is divided into 180 140-km-long blocks.

7.2 The MISR AOD Retrieval Algorithm

In this section, we describe the MISR Level 2 aerosol retrieval algorithm [6]. Let $p =$

Table 1: Examples of single component aerosols, 21 in total.

No.	Name	Radius (μm)	SSA (558nm, Green band)
1	spherical_nonabsorbing (sulfate, sea salt, organic)	0.0562	1
11	spherical_absorbing (sea salt, organic)	1.2835	0.899
19	grains_model1 (dust)	0.7542	0.977

$1, \dots, P$ denote the P pixels on a two-dimensional spatial lattice, i.e. a MISR block. For every pixel p , MISR observes the top-of-atmosphere sun reflectance L_p . Each L_p is a $C = 36$ dimensional vector, corresponding to MISR observations for pixel p at the 36 parallel instrument channels: the four spectral bands in each of the nine viewing angles. Every channel c is weighted by σ_c , $c = 1, \dots, 36$, to account for different measurement or modeling errors. In MISR retrieval algorithm, σ_c is assigned as, $\sigma_c = 0.05 \times \max(\bar{L}_c, 0.04)$, where $\bar{L}_c$ is the average of observed sun reflectances at channel c over all pixels.

Based on a balance between computation expense and current knowledge on aerosols mixings in various geographical locations, MISR retrieval algorithm only considers aerosol compositions containing $M = 3$ or fewer component aerosols, examples shown in Table 2. Our model will extend to allow aerosol compositions containing $M = 4$ component aerosols. For example, the four components involved in aerosol retrieval for the greater Beijing area include spherical non-absorbing aerosols (without sulfate), spherical non-absorbing aerosols (with sulfate), spherical absorbing aerosols, and grains(dust).

For the p th pixel on the lattice, use τ_p to denote its AOD value, and $\theta_p = (\theta_{p1}, \dots, \theta_{pM})$ for the aerosol composition, i.e. values of the associated M -dimensional vector indicating

Table 2: Examples of aerosol compositions, 74 in total

No.	Name	Mixing Percentages of Components (558nm)	SSA (558nm)
5	spherical_nonabsorbing	1: 70%, 5: 30%	1
51	spherical_med_dust	2: 72%, 6: 8%, 19: 20%	0.995
70	spherical_med_coarse_dust	2: 20%, 19: 16%, 21: 64%	0.934

the mixing percentages of the M component aerosols involved. For each p , $\sum_{m=1}^M \theta_{pm} = 1$. MISR retrieval algorithm produces retrievals of AOD τ_p and compositions θ_p based on observed L_p , independently for each pixel p on the spatial lattice, $p = 1, \dots, P$.

Let T denote a pre-fixed grid of AOD values and $s = 1, \dots, 21$ denote the 21 MISR-designated single component aerosols. The Simulated MISR Ancillary Radiative Transfer (SMART) dataset contains the components $\{L^{RT,s}(\tau)\}_{s=1}^{21}$, $\tau \in T$ of radiation fields generated by performing Radiative Transfer (RT) calculations on stratified atmospheric models containing each of the 21 component aerosols with AOD τ [7]. Here we abuse the notation by using M to denote the set of the M chosen component aerosols. Thus, for pixel p with a particular combination of its AOD value $\tau_p \in T$ and compositions θ_p for M , define $L^{RT}(\tau_p, \theta_p)$ as the model top-of-atmosphere sun reflectance:

$$L^{RT}(\tau_p, \theta_p) = \sum_{s \in M} \theta_{ps} L^{RT,s}(\tau_p),$$

a weighted linear combination of the radiation fields $L^{RT,s}(\cdot)$ obtained from SMART dataset.

In order to retrieve AOD τ_p and compositions θ_p for pixel p , MISR retrieval algorithm¹³ first cross-generates a set of combinations of AOD values from a discrete grid and mixing percentages of component aerosols from the 74 MISR pre-fixed choices [6], namely $\theta_p \in \{\theta_0^1, \dots, \theta_0^{74}\}$. Using as input each of the above cross-generated combinations of AOD and aerosol compositions, a Radiative-Transfer-Equation (RTE)-based simulation is conducted to calculate sun reflectance numerically. MISR's standard AOD retrieval algorithm is based on the comparison between such simulated radiances L^{RT} and MISR observations $L_p = (L_{1p}, \dots, L_{Cp})$ on C channels, with their goodness-of-fit measured by [6],

$$\chi_p^2 = \sum_{c=1}^C \frac{(L_{cp} - L_c^{RT}(\tau_p, \theta_p))^2}{2\sigma_c^2}, \quad (8)$$

¹³The model and algorithms in this study focus especially on aerosol retrievals over heterogenous land; retrieval over dark water is conceptually different and should be studied separately.

The combinations of AOD and compositions whose goodness-of-fit measures satisfy the pre-established threshold are considered successful retrievals and contribute to making inference of AOD values.

7.3 Prior Distribution for AOD

We model AOD's spatial dependence structure among adjacent pixels via a intrinsic GMRF prior. Define κ as the homogenous scaler precision and $\mathbf{Q} > 0$ as the precision matrix. Assuming no bias in AOD's smoothness, $\boldsymbol{\tau}$ has its density in the form [23],

$$p(\boldsymbol{\tau}|\kappa, \mathbf{Q}) = (2\pi)^{-\frac{P}{2}} |\kappa \mathbf{Q}|^{\frac{1}{2}} \exp\left(-\frac{\kappa}{2} \boldsymbol{\tau}^\top \mathbf{Q} \boldsymbol{\tau}\right). \quad (9)$$

The precision matrix $\mathbf{Q}$ is symmetric and has the important property that, for pixels $p' \neq p$,

$$\tau_{p'} \perp \tau_p \mid \boldsymbol{\tau}_{-p'p} \iff Q_{p'p} = 0.$$

In our model, for each pixel p , its AOD is directly related only to its few adjacent neighborhoods, suggesting a sparse precision matrix $\mathbf{Q}$. Moreover, employing the intrinsic GMRF of first order [23], i.e. $\mathbf{Q}\mathbf{1} = 0$, the density (9) is invariant to perturbation by the same constant to AOD of all pixels [3],

$$p(\boldsymbol{\tau}|\kappa) \propto \kappa^{\frac{P-1}{2}} \exp\left\{-\frac{\kappa}{2} \sum_{p' \sim p} (\tau_{p'} - \tau_p)^2\right\},$$

where $\sim$ indicates adjacency on the two-dimensional lattice. In this case, the explicit form of precision matrix $\mathbf{Q}$ is,

$$Q_{p'p} = \begin{cases} n_p, & \text{if } p' = p, \\ -1, & \text{if } p' \sim p, \\ 0, & \text{otherwise,} \end{cases}$$

where n_p is the number of adjacent pixels to pixel p ; $n_p = 4$ if all its neighboring pixels are cloudless. This allows us to model the AOD's smoothness regardless of AOD's general level. The full conditional GMRF distribution of τ is,

$$p(\tau_p | \tau_{-p}, \kappa) = \left(\frac{n_p \kappa}{2\pi}\right)^{1/2} \exp \left\{ -\frac{n_p \kappa}{2} \left(\tau_p - \frac{1}{n_p} \sum_{p': p' \sim p} \tau_{p'} \right)^2 \right\},$$

where τ_{-p} are AOD values of all pixels other than the p th.

7.4 Interpolation from SMART

For each of the 21 component aerosols, denoted by $s = 1, \dots, 21$, MISR SMART dataset contains its RT-calculated radiances $\mathbf{L}^{RT,s}$ corresponding to AOD values on a fixed grid, i.e. (0, 0.05, 0.1, 0.2, 0.4, 0.6, 0.8, 1, 1.5, 2, 3, 4, 6). In order to obtain $\mathbf{L}^{RT,s}$ for an arbitrary AOD value between 0 and 6, we consider $\mathbf{L}^{RT,s}$ as a function of AOD τ and interpolate. Examples of $\mathbf{L}^{RT,s}(\tau)$ are shown in Figure 13, for Camera DF and blue band, Camera AN and blue band, Camera DF and red band respectively. An extensive exploratory data analysis suggests a combination of linear and quadratic interpolations to balance between performance and computational efficiency. In particular, linear interpolations are used for $\tau \in (0, 0.4) \cup (3, 6)$, while quadratic interpolations are used for $\tau \in (0.4, 3)$.

7.5 Sensitivity and Computational Expenses of Our MCMC Retrieval Algorithms

We test the stability of outputs from the MCMC retrieval algorithm, using different initial values for the hyperparameters. As an example, results from two retrievals with initial values for $\kappa = 100$ and 200 are plotted in a scatterplot in the left panel of Figure 14. In the right panel plots retrievals with different initial $\alpha = (0.8, 0.4, 0.1, 0.1)$ against $(0.4, 0.4, 0.1, 0.1)$. As is shown in Figure 14, the majority of the points scatter along the 45 degree line with few exceptions, indicating good stability behaviors of our algorithm outputs.

The computation time of the three proposed algorithms are shown in Table 3, which




Figure 13: Examples of single-scattering radiances as a function of AOD, captured by different cameras and in different spectral bands.




Figure 14: Model sensitivity studies for κ and α .

are all implemented on a server with 62.7 GB RAM using a typical MISR radiance imagery. This case shows our MCMC algorithms in general improve the spatial resolution by a factor of 16. Though the parallel MCMC algorithm improves the MCMC algorithm's computational speed by a factor of four, it is still not efficient for online operation. Therefore, a Maximum a Posteriori (MAP) approach based on optimization is developed for operational purpose¹⁴.

Algorithm	Computation Time for Retrieval
MISR (17.6 km)	~ 16.7 minutes (1004.03 sec)
Global MCMC (4.4 km)	~ 6.2 hours (22452.26 sec)
Parallel MCMC (4.4 km)	~ 1.6 hours (5863.77 sec)
MAP (4.4 km)	~ 6 minutes (378.91 sec)

Table 3: Comparison of computation time.

7.6 Specification of Posteriors and Proposal Distributions in the Metropolis-Hastings

To sample τ_p ($p = 1, \dots, P$) in the first P steps of the Gibbs sampler, we need to construct Markov chains with a stationary distribution as τ_p 's univariate marginal posterior distribution:

$$p(\tau_p | \boldsymbol{\tau}_{-p}, \boldsymbol{\theta}, \boldsymbol{\sigma}, \kappa, \mathbf{L}) \propto \exp \left\{ -\frac{n_p \kappa}{2} \left(\tau_p - \frac{1}{n_p} \sum_{p': p' \sim p} \tau_{p'} \right)^2 - \sum_{c=1}^C \frac{(L_{cp} - L_c^{RT}(\tau_p, \boldsymbol{\theta}_p))^2}{2\sigma_c^2} \right\}.$$

We choose the following Gaussian proposal distribution,

$$q(\tau^* | \boldsymbol{\tau}_{-p}, \kappa) \propto \exp \left(-\frac{n_p \kappa}{2} \left(\tau^* - \frac{1}{n_p} \sum_{p': p' \sim p} \tau_{p'} \right)^2 \right),$$

¹⁴For the coherent structure of this study, the MAP retrieval algorithm is not included in this paper.

which is close to the marginal posterior distribution and easy to sample from [12]. After sampling τ^* from $q(\tau^*|\tau_{-p}, \kappa)$, the transition probability from iteration t to $t + 1$ is given by,

$$\tau_p^{(t+1)} = \begin{cases} \tau^*, & \text{w.p. } \min\left(1, \frac{p(\tau^*|\tau_{-p}, \theta, \sigma, \kappa, L)q(\tau_p^{(t)}|\tau_{-p}, \kappa)}{p(\tau_p^{(t)}|\tau_{-p}, \theta, \sigma, \kappa, L)q(\tau^*|\tau_{-p}, \kappa)}\right), \\ \tau_p^{(t)}, & \text{otherwise.} \end{cases}$$

When conditioning on α , τ , and L , the compositions θ_p for each pixel p are independent and can be sampled separately. A Metropolis-Hastings algorithm is implemented with a Dirichlet proposal distribution: $p(\theta_p^*|\alpha) \propto \prod_{m=1}^M (\theta_{pm}^*)^{\alpha_m-1}$.

7.7 Case Study for Improving Retrieval Resolution

We take a closer look at a particular pixel and three of its adjacent retrievable pixels. Table 4 shows the pixel for which our retrieval algorithm and MISR AOD retrievals disagree largely, with one of the largest errors. The center pixel is highlighted in gray, and is chosen due to the AERONET Beijing site within. Our Bayesian AOD retrieval is performed at 4.4 km resolution; both retrievals are based on MISR observation over Beijing area (Location

Table 4: Local comparison of MISR and our model

Location	AERONET	Our Retrievals	MISR
1	0.2060	0.2168	0.1540
2	N/A	0.1870	0.1540
3	N/A	0.1389	0.1540
4	N/A	0.1710	0.1540

Path 122 Block 59) on January 26, 2009. While these four locations are represented by one pixel in MISR AOD algorithm at 17.6 km resolution and thus share the same retrieval output, our statistical retrieval algorithm at 4.4 km resolution produces four different AOD estimates respectively.

Due to the existence of pixel 3, the special features of pixel 1 are averaged away with the relatively high AOD level, which could be one explanation as to why MISR's retrieval is

much lower than AERONET measurement in this particular example. This is one example of the benefits of including spatial smoothness into the model for a closer look at AOD spatial distribution. Certainly, such smoothness depends largely on the scale of observation and retrievals. Our algorithm will fail if certain events influence the local AOD level on a scale smaller than 4.4 km, which is not very uncommon. In that case, an algorithm that retrieves AOD at an even finer resolution is required to capture and model such impacts.

7.8 More Details on Including Spatial Smoothness




Figure 15: AOD retrievals with (top) and without (bottom) spatial smoothness.

Introducing spatial smoothness into the retrieval model is seeking for a balance between particle movements in the open air and existence of local aerosol emissions. The exact calibration for such smoothness requires more studies on the speed and range of particulate matters' movements. However, Figure 15 comparing AOD retrievals with and without spatial smoothness, reveals the lack of an overall spatial structure leads to many cases of very sharp changes between adjacent pixels. Since such abrupt changes is not a very reasonable situation at this level of spatial resolution, the difference between the panels in

Figure 15 is possible results from lack of control for modeling and sampling errors, and confirms our argument for including spatial smoothness. In addition to the errors, retrieval results with a low level of smoothness are also confusing when trying to identify any major patterns of aerosol spatial distribution.

References

- [1] Michelle L. Bell and Devra L. Davis, *Reassessment of the lethal london fog of 1952: Novel indicators of acute and chronic consequences of acute exposure to air pollution*, Environmental Health Perspectives **109** (2001), no. 3, 389–394.
- [2] D. Bergametti, A. L. Dutot, P. BUAT-MENARD, R. LOSNO, and E. REMOUDAKI, *Seasonal variability of the elemental composition of atmospheric aerosol particles over the northwestern mediterranean*. Tellus B **41B** (1989), no. 3, 353–361.
- [3] Julian Besag, *Spatial interaction and the statistical analysis of lattice systems*, Journal of the Royal Statistical Society **B** (1974), 192–236.
- [4] J. A. De Gouw, A. M. Middlebrook, C. Warneke, R. Ahmadov, E. L. Atlas, R. Bahreini, D. R. Blake, C. A. Brock, J. Brioude, D. W. Fahey, F. C. Fensenfeld, J. S. Holloway, M. Le Henaff, R. A. Lueb, Mckeen S. A. J. F. Meagher, D. M. Murphy, C. Paris, D. D. Parrish, A. E. Perring, I. B. Pollack, A. R. Ravishankara, A. L. Robinson, T. B. Ryerson, J. P. Schwarz, Spackman J. R. A. Srinivasan, and L. A. Watts, *Organic aerosol formation downwind from the deepwater horizon oil spill*, Science **331** (2011), no. 6022, 1295–1299.
- [5] D. J. Diner, J. C. Beckert, G. W. Bothwell, and J. I. Rodriguez, *Performance of the misr instrument during its first 20 months in earth orbit*, IEEE Transactions on Geoscience and Remote Sensing **40** (2002), no. 7, 1449–1466.
- [6] David J. Diner, Wedad A. Abdou, Thomas P. Ackerman, Kathleen Crean, Howard R. Gordon, et al. *Level 2 aerosol retrieval algorithm theoretical basis*, JPL **D** (2008), no. 11400.
- [7] David J. Diner, Wedad A. Abdou, Howard R. Gordon, Ralph A. Kahn, Yuri Knyazikhin, John V. Martonchik, Duncan McDonald, Stuart McMuldroch, Ranga Myneni, and Robert A. West, *Level 2 ancillary products and datasets algorithm theoretical basis*, JPL **D** (1999), no. 13402.

- [8] David J. Diner, Jewel C. Beckert, Terrence H. Reilly, Carol J. Bruegge, James E. Conel, Ralph A. Kahn, and John V. Martonchik, *Multi-angle imaging spectroradiometer (misr) - instrument description and experiment overview*, IEEE Transactions on Geoscience and Remote Sensing **36** (1998), no. 4, 1072–1087.
- [9] A. Gelman, G. O. Roberts, and W. R. Gilks, *Efficient metropolis jumping rules*, Bayesian Statistics **5** (1996), 599–607.
- [10] Andrew Gelman and Donald B. Rubin, *Inference from iterative simulation using multiple sequences*, Statistical Science **7** (1992), no. 4, 457–472.
- [11] S. Geman and D. Geman, *Stochastic relaxation, gibbs distributions and the bayesian restoration of images*, IEEE Transactions on Pattern Analysis and Machine Intelligence **6** (1984), no. 6, 721–741.
- [12] John Geweke and Hisashi Tanizaki, *Bayesian estimation of state-space models using the metropolis-hastings algorithm within gibbs sampling*, Computational Statistics and Data Analysis **37** (2001), 151–170.
- [13] David M. Giles, <http://aeronet.gsfc.nasa.gov/>, 2011.
- [14] Jianbin Guo and Yu Chen, *Patterns of seasonal change in air pollution and suggestions on pollution control in beijing*, Ecology and Environmental Sciences **18** (2009), no. 3, 952–956.
- [15] Jiming Hao and Litao Wang, *Improving urban air quality in china: Beijing case study*, Journal of Air and Waste Management Association **55** (2005), no. 9, 1298–1305.
- [16] B. N. Holben, T. F. Eck, Slutsker I. D. Tanré, J. P. Buis, Setzer A. Vermote E. et al. *Aeronet-a federated instrument network and data archive for aerosol characterization*, Remote Sens. Environ. **66** (1998), no. 1, 1–16.
- [17] Xin Jiang, Yang Liu, Bin Yu, and Ming Jiang, *Comparison of misr aerosol optical thickness with aeronet measurements in beijing metropolitan area*, Remote Sensing of Environment **107** (2007), 45–53.
- [18] Ralph A. Kahn, Yang Chen, David L. Nelson, Fok-Yan Leung, Qinbin Li, David J. Diner, and Jennifer A. Logan, *Wild smoke injection heights: Two perspectives from space*, Geophysical Research Letters **35** (2008), L04809.
- [19] C. D. Koven and I. Fung, *Identifying global dust source areas using high-resolution land surface form*, Journal of Geophysical Research Atmospheres **113** (2008), D22204.

- [20] Kuo-nan Liou, *An introduction to atmospheric radiation*, International Geophysics Series, Academic Press, 2002.
- [21] Marjorie Monleau, Cyrill Bussy, Philippe Lestaevel, Pascale Houpert, François Paquet, and Valérie Chazel, *Bioaccumulation and behavioural effects of depleted uranium in rats exposed to repeated inhalations*, *Neuroscience Letters* **390** (2005), 31–36.
- [22] C. Arden Pope III, Richard T. Burnett, Michael J. Thun, Eugenia E. Calle, Daniel Krewski, Kazuhiko Ito, and George D. Thurston, *Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution*, *Journal of the American Medical Association* **287** (2002), no. 9, 1132–1141.
- [23] Havard Rue and Leonhard Held, *Gaussian markov random fields: Theory and applications*, Chapman and Hall / CRC, 2005.
- [24] F. Solmon, F. Giorgi, and C. Liousse, *Aerosol modelling for regional climate studies: application to anthropogenic particles and evaluation over a european/african domain*, *Tellus* **58B** (2006), 51–72.
- [25] David G. Streets, Joshua S. Fu, Carey J. Jang, Jiming Hao, Kebin He, Xiaoyan Tang, Yuanhang Zhang, Zifa Wang, Zuopan Li, Qiang Zhang, Litao Wang, Binyu Wang, and Carolyne Yu, *Air quality during the 2008 beijing olympic games*, *Atmospheric Environment* **41** (2007), 480–492.
- [26] John W. Winchester and Gordon D. Nifong, *Water pollution in lake michigan by trace elements from pollution aerosol fallout*, *Water, Air, and Soil Pollution* **1** (1971), no. 1, 50–64.